

A Geometric Description of Equivariant K-Homology for Proper Actions

Paul Baum, Nigel Higson and Thomas Schick

ABSTRACT. Let G be a discrete group and let X be a G -finite, proper G -CW-complex. We prove that Kasparov's equivariant K-homology groups $KK_*^G(C_0(X), \mathbb{C})$ are isomorphic to the geometric equivariant K-homology groups of X that are obtained by making the geometric K-homology theory of Baum and Douglas equivariant in the natural way. This reconciles the original and current formulations of the Baum-Connes conjecture for discrete groups.

*Dedicated with admiration and affection to
Alain Connes on his 60th birthday.*

1. Introduction

In the first formulation of the Baum-Connes conjecture [BC00], the topological K-theory of a discrete group G (the “left-hand side” of the conjecture) was defined geometrically in terms of proper G -manifolds. Later, in [BCH94], the definition was changed so as to involve Kasparov's equivariant KK-theory. The change was made to accommodate new examples beyond the realm of discrete groups, such as p -adic groups, for which the geometric definition was not convenient or adequate. But it left open the question of whether the original and revised definitions are equivalent for discrete groups. This is the question that we shall address in this paper.

In a recent article [BHS07], we gave a complete proof that the (non-equivariant) geometric K-homology theory of Baum and Douglas [BD82] agrees with Kasparov's K-homology on finite CW-complexes. Here we shall show that our techniques extend to show that the original and revised definitions of topological K-theory for a discrete group agree—provided

NH and PB were partially supported by grants from the US National Science Foundation. TS was supported by the Courant Research Center “Higher order structures in mathematics” via the German Initiative of Excellence.

that those techniques are supplemented by a key result of Lück and Oliver about equivariant vector bundles over G -finite, proper G -CW-complexes [L001].

Lück and Oliver prove that if X is a G -finite, proper G -CW-complex, then there is a rich supply of equivariant vector bundles on X . It follows that the Grothendieck group $K_G^0(X)$ of complex G -vector bundles is the degree zero group of a $\mathbb{Z}/2\mathbb{Z}$ -graded cohomology theory on X , and it is essentially this fact that we shall need to carry over the arguments of [BHS07] to the equivariant case.

We shall show that the Lück-Oliver theorem is equivalent to the assertion that the crossed product C^* -algebra $C^*(X, G)$ associated to the action of G on X has an approximate identity consisting of projections. As a result $K_G^0(X)$ is isomorphic to the K_0 -group of the crossed product C^* -algebra. This has some further benefits for us—for example it makes it clear that each complex G -vector bundles on a G -compact proper G -manifold has a unique smooth structure, up to isomorphism.

Returning to the Baum-Connes conjecture, the assertion that the old and the revised versions are the same is a consequence of the assertion that the natural G -equivariant development of the Baum-Douglas K -homology theory, which we shall write as $K_*^G(X)$, is isomorphic to Kasparov's group $KK_*^G(C_0(X), \mathbb{C})$ for any G -finite, proper G -CW-complex X . There is a natural map

$$\mu: K_*^G(X) \longrightarrow KK_*^G(C_0(X), \mathbb{C})$$

which is defined using the index of Dirac operators, and we shall prove that it is an isomorphism. What makes this nontrivial is that the groups $K_*^G(X)$ do not obviously constitute a homology theory. We shall address this problem by introducing groups $k_*^G(X)$ that manifestly *do* constitute a homology theory and by constructing a commuting diagram

$$\begin{array}{ccc} k_*^G(X) & \longrightarrow & K_*^G(X) \\ \parallel & & \downarrow \mu \\ k_*^G(X) & \xrightarrow{\mu} & KK_*^G(C_0(X), \mathbb{C}). \end{array}$$

We shall prove that the map from $k_*^G(X)$ to $KK_*^G(C_0(X), \mathbb{C})$ is an isomorphism when X is a G -finite, proper G -CW-complex and that map from $k_*^G(X)$ to $K_*^G(X)$ is surjective. This proves that the map from $k_*^G(X)$ to $KK_*^G(C_0(X), \mathbb{C})$ is an isomorphism.

A more general, but less concrete model for equivariant K -theory, based on correspondences, is given by Emerson and Meyer in [EM08]. They describe in the end a bivariant K -theory for spaces with proper groupoid

actions. Their theory does not always coincide with KK-theory, but when their theory overlaps with ours the two coincide.

In [BOOS09] it is shown that for compact Lie groups the result corresponding to the main result in this paper holds (a direct construction is given of an inverse to the transformation μ). The basis of the theorem is again the fact that for compact Lie group actions there are “enough” equivariant vector bundles.

2. Proper Actions

Throughout the paper we shall work with a fixed a countable discrete group G . By a G -space we shall mean a topological space with an action of G by homeomorphisms. We shall be concerned in the first place with proper G -CW-complexes. These are G -spaces with filtrations

$$\emptyset = X^{-1} \subseteq X^0 \subseteq X^1 \subseteq \dots \subseteq X$$

such that X^k is obtained from X^{k-1} by attaching equivariant cells of the form $D^k \times G/H$ along their boundaries, where H is any finite subgroup of G . See [Lüc05, Section 1] for more details.

The Baum-Connes conjecture as formulated in [BCH94] involves *universal proper G -spaces*. In the context of proper G -CW-complexes, these may be characterized as follows:

2.1. THEOREM ([Lüc05, Theorem 1.9]). *There is a proper G -CW-complex $\underline{E}G$ with the property that if Y is any proper G -CW-complex, then there is a G -equivariant continuous map from Y into $\underline{E}G$, and moreover this map is unique up to equivariant homotopy.*

Clearly the G -CW-complex $\underline{E}G$ is unique up to equivariant homotopy. The universal space used in the formulation of the Baum-Connes conjecture is defined a bit differently (see [BCH94, Section 1]), but by the results of [Lüc05, Section 2] the same conjecture results if the above version of $\underline{E}G$ is used. Compare also Theorem 2.3 below.

A proper G -CW-complex is said to be *G -finite* if only finitely many equivariant cells are used in its construction. These are G -compact proper G -spaces in the sense of the following definition.

2.2. DEFINITION. We shall say that a G -space X is a *G -compact, proper G -space* if

- (a) X is locally compact and Hausdorff.
- (b) The quotient space X/G is compact and Hausdorff in the quotient topology.
- (c) Each point of X is contained in an equivariant neighborhood U that maps continuously and equivariantly onto some proper orbit space G/H (where H is a finite subgroup of G).

Apart from G -finite G -CW-complexes, we shall also be concerned with smooth manifolds (with smooth actions of G) that satisfy these conditions. We shall call them *G -compact proper G -manifolds*.

The following result is a consequence of [Lüc05, Theorem 3.7] (the final statement reflects a simple feature of the CW-topology on $\underline{E}G$).

2.3. THEOREM. *If X is any G -finite proper G -space, then there is a G -equivariant map from X to $\underline{E}G$. It is unique up to equivariant homotopy, and its image is contained within a G -finite subcomplex of $\underline{E}G$.*

3. Equivariant Geometric K-Homology

In this section we shall present the equivariant version of the geometric K-homology theory of Baum and Douglas [BD82]. The definition presents no difficulties, so we shall be brief. The reader is referred to [BD82] or [BHS07] for treatments of the non-equivariant theory.

We shall work with principal bundles, rather than with spinor bundles as in [BD82] or [BHS07]. To fix notation, recall the following rudimentary facts about Clifford algebras, Spin^c -groups and Spin^c -structures. Denote by $\text{Cliff}(\mathfrak{n})$ the $\mathbb{Z}/2\mathbb{Z}$ -graded complex $*$ -algebra generated by skew-adjoint degree-one elements e_1, \dots, e_n such that

$$e_i e_j + e_j e_i = -2\delta_{ij}I.$$

We shall consider \mathbb{R}^n as embedded into $\text{Cliff}(\mathfrak{n})$ in such a way that the standard basis of \mathbb{R}^n is carried to e_1, \dots, e_n .

Denote by $\text{Spin}^c(\mathfrak{n})$ the group of all even-grading-degree unitary elements in $\text{Cliff}(\mathfrak{n})$ that map \mathbb{R}^n into itself under the adjoint action. This is a compact Lie group. The image of the group homomorphism

$$\alpha: \text{Spin}^c(\mathfrak{n}) \longrightarrow \text{GL}(\mathfrak{n}, \mathbb{R})$$

given by the adjoint action is $\text{SO}(\mathfrak{n})$ and the kernel is the circle group $\text{U}(1)$ of all unitaries in $\text{Cliff}(\mathfrak{n})$ that are multiples of the identity element.

There is a natural complex conjugation operation on the algebra $\text{Cliff}(\mathfrak{n})$ (since the relations defining the Clifford algebra involve only real coefficients) and the map $u \mapsto u\bar{u}^*$ is a homomorphism from $\text{Spin}^c(\mathfrak{n})$ onto $\text{U}(1)$. The combined homomorphism

$$\text{Spin}^c(\mathfrak{n}) \longrightarrow \text{SO}(\mathfrak{n}) \times \text{U}(1)$$

is a double covering.

Let M be a smooth, proper G -manifold and let V be a smooth, real G -vector bundle over M of rank n . A G - Spin^c -structure on V is a homotopy class of reductions of the principal frame bundle of V (viewed as a

G-equivariant right principal $GL(n, \mathbb{R})$ -bundle) to a G-equivariant principal $Spin^c(n)$ -bundle. In other words it is a G-homotopy class of commuting diagrams

$$\begin{array}{ccc} Q & \xrightarrow{\varphi} & P \\ \downarrow & & \downarrow \\ M & \xlongequal{\quad} & M \end{array}$$

of smooth G-manifolds, where P is the bundle of ordered bases for the fibers of V, Q is a G-equivariant principal $Spin^c(n)$ -bundle, and

$$\varphi(qu) = \varphi(q)\alpha(u)$$

for every $q \in Q$ and every $u \in Spin^c(n)$. A G- $Spin^c$ -structure on V determines a G-invariant orientation, and a specific choice of Q within its homotopy class determines a Euclidean structure.

3.1. EXAMPLE. Every G-equivariant complex vector bundle carries a natural $Spin^c$ -structure because there is a (unique) group homomorphism $U(k) \rightarrow Spin^c(2k)$ that lifts the map

$$U(k) \longrightarrow SO(2k) \times U(1)$$

given in the right-hand factor by the determinant.

A G- $Spin^c$ -vector bundle is a smooth real G-vector bundle with a given G- $Spin^c$ -structure. The direct sum of two G- $Spin^c$ -vector bundles carries a natural G- $Spin^c$ -structure. It is obtained from the diagram

$$\begin{array}{ccc} Spin^c(m) \times Spin^c(n) & \longrightarrow & Spin^c(m+n) \\ \downarrow & & \downarrow \\ GL(n, \mathbb{R}) \times GL(m, \mathbb{R}) & \longrightarrow & GL(m+n, \mathbb{R}) \end{array}$$

that is in turn obtained from the inclusions of $Cliff(m)$ and $Cliff(n)$ into $Cliff(m+n)$ given by the formulas $e_k \mapsto e_k$ and $e_k \mapsto e_{m+k}$, respectively. In addition, if V and $V \oplus W$ carry G- $Spin^c$ -structures, then there is a unique G- $Spin^c$ -structure on W whose direct sum, as above, with the G- $Spin^c$ -structure on V is the given G- $Spin^c$ -structure on the direct sum. This is the *two out of three principle* for $Spin^c$ -structures.

If M is a smooth G-manifold, then a G- $Spin^c$ -structure on M is a G- $Spin^c$ -structure on its tangent bundle, and a G- $Spin^c$ -manifold is a smooth G-manifold together with a given G- $Spin^c$ -structure.

3.2. DEFINITION. Let X be any G-space. An equivariant K-cycle for X is a triple (M, E, f) consisting of:

- (a) A G -compact, proper G - Spin^c -manifold M without boundary.¹
- (b) A smooth complex G -vector bundle E over M .
- (c) A continuous and G -equivariant map $f: M \rightarrow X$.

The geometric equivariant K -homology groups $K_*^G(X)$ will be obtained by placing a certain equivalence relation on the class of all equivariant K -cycles. Before describing it, we give constructions at the level of cycles that will give the arithmetic structure of the groups $K_*^G(X)$.

If (M, E, f) and (M', E', f') are two equivariant K -cycles for X , then their *disjoint union* is the equivariant K -cycle $(M \sqcup M', E \sqcup E', f \sqcup f')$. The operation of disjoint union will give addition.

Let V be a G - Spin^c -vector bundle with a G - Spin^c -structure $\varphi: Q \rightarrow P$. Fix an orientation-reversing isometry of \mathbb{R}^n . Since it preserves the inner product, τ induces an automorphism of $\text{Cliff}(n)$, and hence of $\text{Spin}^c(n)$, that we shall also denote by τ . Consider the map $\varphi_\tau: Q_\tau \rightarrow P$, where:

- (a) Q_τ is equal to Q as a G -manifold, but has the twisted action given by the formula $q \cdot_\tau u = q \cdot \tau(u)$ of the group $\text{Spin}^c(n)$.
- (b) $\phi_\tau(q) = \phi(q)\tau$.

It defines the *opposite* G - Spin^c -vector bundle $-V$. Applying this to manifolds, we define the *opposite* of an equivariant K -cycle (M, E, f) to be the equivariant K -cycle $(-M, E, f)$. This will give the operation of additive inverse in the geometric groups $K_*^G(X)$.

If M is a Spin^c - G -manifold, then its boundary ∂M inherits a Spin^c - G -structure. This is obtained from the pullback diagram

$$\begin{array}{ccc} \text{Spin}^c(n-1) & \longrightarrow & \text{Spin}^c(n) \\ \downarrow & & \downarrow \\ \text{GL}(n-1, \mathbb{R}) & \longrightarrow & \text{GL}(n, \mathbb{R}) \end{array}$$

associated to the lower-right-corner embedding of $\text{GL}(n-1, \mathbb{R})$ into $\text{GL}(n, \mathbb{R})$ and the inclusion of $\text{Cliff}(n-1)$ into $\text{Cliff}(n)$ that maps the generators e_k to e_{k+1} . Using the outward-pointing normal first convention, the bundle of frames for $T\partial M$ maps to the restriction to the boundary of the bundle of frames for TM . A pullback construction gives the required reduction to $\text{Spin}^c(n-1)$.

3.3. DEFINITION. An equivariant K -cycle for X is a *boundary* if there is a G -compact, proper G - Spin^c -manifold W with boundary, a smooth, Hermitian equivariant vector bundle E over W and a continuous equivariant map $f: W \rightarrow X$ such that the given cycle is isomorphic to $(\partial W, E|_{\partial W}, f|_{\partial W})$.

¹The manifold M need not be connected. Moreover different connected components of M may have different dimensions.

Two equivariant K-cycles for X , (M_1, E_1, f_1) and (M_2, E_2, f_2) are *bordant* if the disjoint union of one with the opposite of the other is a boundary.

The most subtle aspect of the equivalence relation on equivariant K-cycles that defines geometric K-homology involves certain sphere bundles over Spin^c -manifolds. To describe it we begin by considering a single sphere.

View S^{n-1} as the boundary of the unit ball in \mathbb{R}^n . The frame bundle for \mathbb{R}^n can of course be identified with $\mathbb{R}^n \times \text{GL}(n, \mathbb{R})$ since the columns of any invertible matrix constitute a frame for \mathbb{R}^n . We can therefore equip \mathbb{R}^n with the *trivial* Spin^c -structure $\mathbb{R}^n \times \text{Spin}^c(n)$.

According to the prescription given prior to Definition 3.3, the associated Spin^c -structure on the sphere S^{n-1} is given by the right principal $\text{Spin}^c(n-1)$ -bundle Q whose fiber at $v \in S^{n-1}$ is the space of all elements $u \in \text{Spin}^c(n)$ whose image in $\text{SO}(n)$ is a matrix with first column equal to v . Observe that Q is $\text{Spin}^c(n)$ -equivariant for the left action of $\text{Spin}^c(n)$ on the sphere given by the projection to $\text{SO}(n)$.

Let us now assume that $n = 2k + 1$. We are going to fix a certain $\text{Spin}^c(n)$ -equivariant complex vector bundle F on S^{2k} . The key property of F is that the $\text{Spin}^c(n)$ -equivariant index of the Dirac operator (discussed in the next section) coupled to F is equal to the rank-one trivial representation of $\text{Spin}^c(n)$. An explicit calculation, given in [BHS07], shows that the dual of the positive part of the spinor bundle for S^{n-1} has the required property. It follows easily from the Bott periodicity theorem that F is essentially unique (up to addition of trivial bundles, any two F are isomorphic). For what follows, any choice of F will do. The bundle F and the trivial line bundle together generate $K(S^{2k})$, and for that reason we shall call it the *Bott generator*.

Following these preliminaries, we can describe the “vector bundle modification” step in the equivalence relation defining geometric K-homology.

Let V be a G - Spin^c -vector bundle of rank $2k$ over a G - Spin^c -manifold M and denote by \widehat{M} the sphere bundle² of the direct sum vector bundle $\mathbb{R} \oplus V$. The manifold \widehat{M} may be described as the fiber bundle

$$\widehat{M} = Q \times_{\text{Spin}^c(2k+1)} S^{2k},$$

where Q is the principal G - $\text{Spin}^c(2k+1)$ -bundle associated to $\mathbb{R} \oplus V$. Its tangent bundle is isomorphic to the pullback of the tangent bundle of M , direct sum the fiberwise tangent bundle $Q \times_{\text{Spin}^c(2k+1)} \text{TS}^{2k}$. Both carry natural G - Spin^c -structures, and so \widehat{M} is a G - Spin^c -manifold.

²Strictly speaking, to form the sphere bundle we need a metric on V and so a specific choice of principal bundle Q within its homotopy class. Of course, any two sphere bundles will be bordant.

Form the G -equivariant complex vector bundle

$$Q \times_{\text{Spin}^c(2k+1)} F,$$

from the Bott generator discussed above. We shall use the same symbol F for this bundle over \widehat{M} .

3.4. DEFINITION. Let (M, E, f) be an equivariant K -cycle and let V be a rank $2k$ G - Spin^c -vector bundle over M . The *modification* of (M, E, f) associated to V is the equivariant K -cycle

$$(M, E, f)^\wedge = (\widehat{M}, F \otimes \pi^*(E), f \circ \pi),$$

where:

- (a) \widehat{M} is the total space of the sphere bundle of $\mathbb{R} \oplus V$, equipped with the G - Spin^c -structure described above;
- (b) π is the projection from \widehat{M} onto M ; and
- (c) F is the G -equivariant complex vector bundle on \widehat{M} described above.

We are now ready to define the geometric equivariant K -homology groups.

3.5. DEFINITION. Denote by $K^G(X)$ the set of equivalence classes of equivariant K -cycles over X , for the equivalence relation generated by the following three elementary relations:

- (a) If (M, E_1, f) and (M, E_2, f) are two equivariant K -cycles with the same proper, G -compact G - Spin^c -manifold M and same $f: M \rightarrow X$, then

$$(M \sqcup M, E_1 \sqcup E_2, f \sqcup f) \sim (M, E_1 \oplus E_2, f).$$

- (b) If (M_1, E_1, f_1) and (M_2, E_2, f_2) are bordant equivariant K -cycles then

$$(M_1, E_1, f_1) \sim (M_2, E_2, f_2).$$

- (c) If (M, E, f) is an equivariant K -cycle, if V is an even-rank G - Spin^c -vector bundle over M , and if $(M, E, f)^\wedge$ is the modification of (M, E, f) associated to V , then

$$(M, E, f) \sim (M, E, f)^\wedge.$$

The set $K^G(X)$ is an abelian group with addition given by disjoint union. Denote by $K_{\text{ev}}^G(X)$ and $K_{\text{odd}}^G(X)$ the subgroups of $K^G(X)$ composed of equivalence classes of equivariant K -cycles (M, E, f) for which every connected component of M is even-dimensional or odd-dimensional, respectively. Then $K^G(X) \cong K_{\text{ev}}^G(X) \oplus K_{\text{odd}}^G(X)$.

4. Equivariant Kasparov Theory

In this section we shall define a natural transformation from geometric equivariant K-homology to Kasparov's equivariant K-homology. Once again, this is a straightforward extension to the equivariant context of the Baum-Douglas theory that was reviewed in detail already in [BHS07]. Because of this we shall be brief.

Fix a *second countable* G -compact proper G -space X , for example a G -finite proper G -CW-complex. The second countability assumption is made for consistency with Kasparov's theory, which applies to second countable locally compact spaces, or separable C^* -algebras.

We shall denote by $\mathrm{KK}_n^G(C_0(X), \mathbb{C})$ Kasparov's bivariant KK-theory group $\mathrm{KK}^G(C_0(X), \mathbb{C}\mathrm{li}\mathrm{f}\mathrm{f}(n))$ (the action of G on $\mathbb{C}\mathrm{li}\mathrm{f}\mathrm{f}(n)$ is trivial). See [Kas88, Section 2]. There are canonical isomorphisms

$$\mathrm{KK}_n^G(C_0(X), \mathbb{C}) \cong \mathrm{KK}_{n+2}^G(C_0(X), \mathbb{C})$$

coming from the periodicity of Clifford algebras. Compare [BHS07]. As a result we may form the 2-periodic groups $\mathrm{KK}_{\mathrm{ev}/\mathrm{odd}}^G(C_0(X), \mathbb{C})$.

The natural transformation

$$\mu: \mathrm{K}_{\mathrm{ev}/\mathrm{odd}}^G(X) \longrightarrow \mathrm{KK}_{\mathrm{ev}/\mathrm{odd}}^G(C_0(X), \mathbb{C})$$

into Kasparov theory is defined by associating to an equivariant K-cycle (M, E, f) a Dirac operator, and then constructing from the Dirac operator a cycle for Kasparov's analytic K-homology group.

The vector space $\mathbb{C}\mathrm{li}\mathrm{f}\mathrm{f}(n)$ carries a natural inner product in which the monomials $e_{i_1} \cdots e_{i_k}$ form an orthonormal basis. If M is a G -compact, proper G - Spin^c -manifold, and if Q is a lifting to $\mathrm{Spin}^c(n)$ of the frame bundle of M , then the $\mathbb{Z}/2\mathbb{Z}$ -graded Hermitian vector bundle

$$S = Q \times_{\mathrm{Spin}(n)} \mathbb{C}\mathrm{li}\mathrm{f}\mathrm{f}(n)$$

that is formed using the left multiplication action of $\mathrm{Spin}^c(n)$ on $\mathbb{C}\mathrm{li}\mathrm{f}\mathrm{f}(n)$ carries a right action of the algebra $\mathbb{C}\mathrm{li}\mathrm{f}\mathrm{f}(n)$ and a commuting left action of TM as odd-graded skew-adjoint endomorphisms such that $v^2 = -\|v\|^2 I$. This is called the action of TM on the *spinor bundle* S by *Clifford multiplication*.

4.1. REMARK. There are other versions of the spinor bundle that do not carry a right Clifford algebra action. The bundle used here has the advantage of allowing a uniform treatment of both even and odd-rank bundles V . In addition the real case may be treated similarly (although we shall not consider it in this paper).

4.2. DEFINITION. Let M be a G -compact proper G - Spin^c -manifold. Fix an associated principal Spin^c -bundle over M , and let S be the spinor

bundle, as above. Let E be a smooth, Hermitian G -vector bundle over M . We shall call an odd-graded, symmetric, first-order G -equivariant linear partial differential operator D acting on the sections of $S \otimes E$ a *Dirac operator* if it commutes with the right Clifford algebra action on the spinor bundle and if

$$[D, f]u = \text{grad}(f) \cdot u,$$

for every smooth function f on M and every section u of $S \otimes E$, where $\text{grad}(f) \cdot u$ denotes Clifford multiplication on S by the gradient of f .

Dirac operators in this sense always exist, and basic PDE theory gives the following result:

4.3. PROPOSITION. *The Dirac operator D , considered as an unbounded operator on $L^2(M, S \otimes E)$ with domain the smooth compactly supported sections is essentially self-adjoint. The bounded G -equivariant Hilbert space operator $F = D(I + D^2)^{-1/2}$ commutes, modulo compact operators, with multiplication operators from $C_0(M)$. Moreover the product of $I - F^2$ with any multiplication operator from $C_0(M)$ is a compact operator. \square*

Now the Hilbert space $L^2(M, S \otimes E)$ carries a right action of $\text{Cliff}(\mathfrak{n})$ that commutes with D and the action of $C_0(M)$. It also carries a unique $\text{Cliff}(\mathfrak{n})$ -valued inner product $\langle \cdot, \cdot \rangle_{\text{Cliff}}$ such that

$$\langle s_1, s_2 \rangle = \tau(\langle s_1, s_2 \rangle_{\text{Cliff}})$$

where on the left is the L^2 -inner product, and on the right is the state τ on $\text{Cliff}(\mathfrak{n})$ that maps all nontrivial monomials $e_{i_1} \cdots e_{i_p}$ to zero. Using it we place a Hilbert $\text{Cliff}(\mathfrak{n})$ -module structure on $L^2(M, S \otimes E)$.

Proposition 4.3 implies that the operator $F = D(I + D^2)^{-1/2}$, viewed as an operator on the Hilbert $\text{Cliff}(\mathfrak{n})$ -module $L^2(M, S \otimes E)$, yields a cycle for Kasparov's equivariant KK -group $\text{KK}^G(C_0(M), \text{Cliff}(\mathfrak{n}))$ (see [Kas88, Definition 2.2]).

4.4. DEFINITION. We shall denote by $[M, E] \in \text{KK}_n^G(C_0(M), \mathbb{C})$ the KK -class of the operator $F = D(I + D^2)^{-1/2}$.

The first main theorem concerning the classes $[M, E]$ is as follows:

4.5. THEOREM. *The correspondence that associates to each equivariant K -cycle (M, E, f) the KK -class*

$$f_*[M, E] \in \text{KK}_{\text{ev}/\text{odd}}^G(C_0(X), \mathbb{C})$$

gives a well-defined homomorphism

$$\mu: \text{K}_{\text{ev}/\text{odd}}^G(X) \longrightarrow \text{KK}_{\text{ev}/\text{odd}}^G(C_0(X), \mathbb{C}).$$

The non-equivariant case of the theorem is proved in [BHS07]. The proof for the equivariant case is exactly the same and therefore will not be repeated.

Our aim in this paper is to prove the second main theorem concerning the classes $[M, E]$.

4.6. THEOREM. *If X is any proper, G -finite G -CW-complex, then the index map*

$$\mu: K_{\text{ev/odd}}^G(X) \longrightarrow KK_{\text{ev/odd}}^G(C_0(X), \mathbb{C})$$

is an isomorphism.

The non-equivariant version of the theorem is due to Baum and Douglas, and is proved in detail in [BHS07]. Although the proof of the equivariant result is the same in outline, new issues must also be resolved having to do with the properties of equivariant vector bundles on G -compact proper G -spaces. These we shall consider next.

5. Equivariant Vector Bundles

Throughout this section we shall use the term G -bundle as an abbreviation for G -equivariant complex vector bundle. We shall review the basic theory of G -bundles over G -compact proper G -spaces, mostly as worked out by Lück and Oliver in [LO01]. In the next section we shall recast their results in the language of C^* -algebra K -theory.

5.1. THEOREM. *Let X be a G -compact, proper G -space. There is a G -bundle E over X such that for every $x \in X$, the fiber E_x is contained in a multiple of the regular representation of the isotropy group G_x . \square*

PROOF. This is proved for G -finite proper G -CW-complexes in [LO01, Corollary 2.8]. That result extends to more general X by pulling back along the map supplied by Theorem 2.3. \square

5.2. COROLLARY (Compare [LO01, Lemma 3.8]). *Let Z be a G -compact proper G -space and let X be a closed, G -invariant subset of Z . If F is any G -bundle on X , then there is a G -bundle E on Z such that F embeds as a summand of $E|_X$.*

PROOF. Fix a G -bundle E on Z , as in Theorem 5.1. There are G -invariant open subsets U_1, \dots, U_n of X such that:

- (a) The sets cover X .
- (b) For each j there is a finite subgroup of $F_j \subseteq G$ and an equivariant map $\pi_j: U_j \rightarrow G/F_j$.
- (c) $F|_{U_j}$ is isomorphic to a bundle pulled back along π_j .
- (d) $E|_{U_j}$ is also isomorphic to a bundle pulled back along π_j .

Replacing E by a direct sum $E \oplus \cdots \oplus E$, if necessary, we find that $F|_{U_j}$ may be embedded as a summand of $E|_{U_j}$, for every j . Making a second replacement of E by an n -fold direct sum $E \oplus \cdots \oplus E$ and using a standard partition of unity argument, we may now embed F into E , as required. \square

More generally, if $f: X \rightarrow Z$ is a map between G -compact proper G -spaces, and if F is a G -bundle on X , then the same argument shows that F is isomorphic to a summand of the pullback along f of some G -bundle on Z .

5.3. DEFINITION. If S is any set, then denote by $\mathbb{C}[S]$ the free vector space on the set S , equipped with the standard inner product in which the elements of S are orthonormal. If S is equipped with an action of G , then we shall consider $\mathbb{C}[S]$ to be equipped with the corresponding permutation action of G .

We are interested primarily in the case where $S = G$, which we shall view as equipped with the usual left translation action of G .

5.4. DEFINITION. A *standard G -bundle* on a G -compact proper G -space X is a G -invariant subset E of $X \times \mathbb{C}[G]$ with the property that for every compact subset $K \subseteq X$ there is a *finite* subset $S \subseteq G$ such that the intersection of E with $K \times \mathbb{C}[G]$ is a (nonequivariant) complex vector subbundle of $K \times \mathbb{C}[S]$.

5.5. REMARK. We require that the restriction of E to K , as above, be a topological vector subbundle of the finite-dimensional trivial bundle $K \times \mathbb{C}[S]$. This fixes the topology on E and determines a G -bundle structure.

It follows from a standard partition of unity argument that every G -bundle on X is isomorphic to a standard G -bundle. We are going to prove the following result, which gives the set of standard G -bundles a useful directed set structure.

5.6. THEOREM. *Any two standard G -bundles are subbundles of a common third. Moreover the union of all standard G -bundles is $X \times \mathbb{C}[G]$.*

5.7. REMARK. In Section 8 we shall modify Definition 5.4 very slightly by replacing G with a countable disjoint union $G_\infty = G \sqcup G \sqcup \cdots$ (thought of as a left G -set). Theorem 5.6 remains true, with the same proof.

Since the theorem is obvious if G is finite, we shall assume G is infinite until the proof of Theorem 5.6 is concluded.

5.8. LEMMA. *If S is a finite subset of G , if K is a compact subset of X , and if E is any G -bundle over X , then there is a standard G -bundle E_1 that is isomorphic to E and whose restriction to K is orthogonal to $K \times \mathbb{C}[S]$.*

PROOF. Let E_1 be any standard bundle that is isomorphic to the equivariant vector bundle E . If $g \in G$, then set

$$E_1 \cdot g = \{ (x, e \cdot g) : e \in E_{1,x} \}$$

is also standard G -bundle. Here, in forming the vectors $e \cdot g$ we are using the *right* translation action of G on itself and hence on $\mathbb{C}[G]$. The bundle $E_1 \cdot g$ is isomorphic to E_1 , and hence to E . If we enlarge S , if necessary, so that $E_1|_K \subseteq K \times \mathbb{C}[S]$, and if we choose $g \in G$ so that $S \cap Sg = \emptyset$, then $(E_1 \cdot g)|_K$ is orthogonal to $K \times \mathbb{C}[S]$, as required. \square

5.9. LEMMA. *Let E be a G -bundle on X and let E_2 be a standard G -bundle. There is a standard G -bundle that is isomorphic to E and orthogonal to E_2 .*

PROOF. Let K be a compact subset of X whose G -saturation is X , and let S be a finite subset of G such that $E_2|_K \subseteq K \times \mathbb{C}[S]$. Now apply the previous lemma. \square

5.10. LEMMA. *Let U be a G -invariant open subset of X , and let Y and Z be G -invariant closed subsets of X such that*

$$Z \subseteq U \subseteq Y \subseteq X.$$

Let F be a standard G -bundle over Y . There is a standard G -bundle E_1 over X such that $F|_Z \subseteq E_1|_Z$. Moreover given a standard G -bundle E_2 over X such that $E_2|_Y$ is orthogonal to F , the standard G -bundle E_1 may be chosen to be orthogonal to E_2 .

PROOF. According to Corollary 5.2, there is a G -bundle E over X such that F embeds in $E|_Y$. Any complement of the image of F in $E|_Y$ may be embedded as a standard G -bundle F' on Y that is orthogonal to $F \oplus E_2|_Y$.

We can choose an isomorphism $\Phi: E|_Y \rightarrow F \oplus F'$. Next, there is an embedding Ψ of E as a standard G -bundle on X such that $\Psi[E]$ is orthogonal to E_2 and $\Psi[E]|_Y$ is orthogonal to F (by a slight elaboration of Lemma 5.9). If we choose a G -invariant scalar function φ on X such that $\varphi = 1$ on Z and $\varphi = 0$ outside of U , and if we set $\psi = 1 - \varphi$, then $E_1 = (\varphi\Phi + \psi\Psi)[E]$ has the required properties. \square

5.11. LEMMA. *Let K be any compact subset of X , and let S be a finite subset of G . There is a standard G -bundle that contains $K \times \mathbb{C}[S]$.*

PROOF. The compact set K may be written as a finite union of compact sets

$$K = K_1 \cup \cdots \cup K_n$$

where each K_j is included in a G -invariant open set that maps equivariantly onto some proper coset space G/H_j , in such a way that K_j maps to the identity coset. We shall use induction on n .

Let E_2 be a standard G -bundle that contains the set

$$(K_1 \cup \cdots \cup K_{n-1}) \times \mathbb{C}[S].$$

There is a G -compact subset $Y \subseteq X$ that contains a G -invariant neighborhood U of K_n and over which there is a standard G -bundle L such that

$$K_n \times \mathbb{C}[S] \subseteq L|_{K_n} \quad \text{and} \quad E_2|_Y \subseteq L.$$

Indeed we may choose Y so that it maps equivariantly to G/H_n , and if Y_n is the inverse image of the identity coset, then we may form

$$L = \bigcup_{g \in G} gY_n \times \mathbb{C}[S_n g],$$

where S_n is a sufficiently large finite and right H_n -invariant subset of G .

Now apply the previous lemma to the standard G -bundle $F = L \ominus E_2|_Y$ (the orthogonal complement of $E_2|_Y$ in L) to obtain a standard G -bundle E_1 on X such that E_1 is orthogonal to E_2 and

$$L|_{K_n} \ominus E_2|_{K_n} \subseteq E_1|_{K_n}.$$

The standard G -bundle $E_1 \oplus E_2$ then contains $K \times \mathbb{C}[S]$, as required. \square

PROOF OF THEOREM 5.6. Since there is a standard G -bundle that contains any given $K \times \mathbb{C}[S]$, it is clear that the union of all standard G -bundles is $X \times \mathbb{C}[G]$. Let E_1 and E_2 be standard G -bundles on X . Choose a compact set K whose G -translates cover X and choose a finite set $S \subseteq G$ such that the $E_1|_K, E_2|_K \subseteq K \times \mathbb{C}[S]$. If E is a standard G -bundle containing $K \times \mathbb{C}[S]$, then it contains E_1 and E_2 . \square

6. C^* -Algebras and Equivariant K -Theory

6.1. **DEFINITION.** If S is any set, then denote by $M[S]$ the $*$ -algebra of complex matrices $[T_{s_1 s_2}]$ with rows and columns parametrized by the set S , all but finitely many of whose entries are zero.

We shall be interested in the case where $S = G$. In this case the group G acts on $M[G]$ by automorphisms via the formula $(g \cdot T)_{g_1, g_2} = T_{g^{-1}g_1, g^{-1}g_2}$.

6.2. **DEFINITION.** Let X be a G -compact proper G -space. Let us call a function $F: X \rightarrow M[G]$ *standard* if its matrix element functions

$$F_{g_1, g_2}: x \mapsto F(x)_{g_1, g_2}$$

are continuous and compactly supported, and if for every compact subset K of X all but finitely many of them vanish outside of K . We shall denote by $\mathcal{C}(X, G)$ the $*$ -algebra of all standard, G -equivariant functions from X to $M[G]$.

Note that if P is a projection in the $*$ -algebra $\mathcal{C}(X, G)$, then the range of P (that is the bundle over X whose fiber of $x \in X$ is the range of the projection operator $P(x)$ in $\mathbb{C}[G]$) is a standard G -bundle in the sense of Section 5. In fact every standard G -bundle is obtained in this way, which explains our interest in $\mathcal{C}(X, G)$. In fact we are even more interested in the following C^* -algebra completion of $\mathcal{C}(X, G)$.

6.3. DEFINITION. Let X be a G -compact proper G -space. Denote by $C^*(X, G)$ the C^* -algebra of G -equivariant, continuous functions from X into the compact operators on $\ell^2(G)$.

6.4. REMARK. The C^* -algebra $C^*(X, G)$ is isomorphic to the crossed product C^* -algebra $C_0(X) \rtimes G$. If E^{g_1, g_2} denotes the matrix with 1 in entry (g_1, g_2) and zero in every other entry, then the formula

$$f[g] \mapsto \sum_h h(f) E^{h, hg}$$

gives an isomorphism from $C_0(X) \rtimes G$ to $C^*(X, G)$, and the formula

$$F \mapsto \sum_{g \in G} F_{e, g}[g]$$

gives its inverse. Since the action of G on X is proper, the maximal and reduced crossed products are equal. Indeed there is a unique C^* -algebra completion of the $*$ -algebra $\mathcal{C}(X, G)$.

6.5. LEMMA. *Assume that G is infinite and X is a G -compact proper G -space. The correspondence between projections in $\mathcal{C}(X, G)$ and their ranges induces bijections among the following sets:*

- (a) *Equivalence classes of projections in $\mathcal{C}(X, G)$.*
- (b) *Equivalence classes of projections in $C^*(X, G)$.*
- (c) *Isomorphism classes of standard G -bundles on X .*
- (d) *Isomorphism classes of hermitian G -bundles on X .*

If X is a G -compact proper G -manifold, then there is in addition a bijection with

- (e) *Isomorphism classes of smooth hermitian G -bundles on X .*

PROOF. Recall that two projections P and Q in a $*$ -algebra are equivalent if and only if there is an element U such that $U^*U = P$ and $UU^* = Q$. The inclusion of $\mathcal{C}(X, G)$ into $C^*(X, G)$ is a simple example of a *holomorphically closed subalgebra*, and as a result the inclusion induces a bijection between the sets in (a) and (b). Compare [Bla98, Sections 3 and 4]. The sets in (a) and (c) are in bijective correspondence virtually by definition.

The sets in (c) and (d) are in bijection thanks to Lemma 5.8, which in particular shows that every G -bundle is isomorphic to a standard bundle, if G is infinite.

If X is a manifold, then the inclusion of the smooth functions in $\mathcal{C}(X, G)$ into $C^*(X, G)$ is also a holomorphically closed subalgebra, and this gives the final part of the lemma since equivalence classes projections in the algebra of smooth functions correspond to isomorphism classes of (the obvious concept of) smooth standard G -bundles. \square

6.6. REMARK. If G is finite, then the lemma remains true if $\mathcal{C}(X, G)$ and $C^*(X, G)$ are replaced by direct limits of matrix algebras over themselves.

6.7. THEOREM. *The C^* -algebra $C^*(X, G)$ has an approximate identity consisting of projections.*

PROOF. We claim that for every finite set of elements F_1, \dots, F_n in $\mathcal{C}(X, G)$ there is a projection P in $\mathcal{C}(X, G)$ such that

$$F_j = PF_j = F_jP$$

for all $j = 1, \dots, n$. Indeed the orthogonal projection onto any standard G -bundle is a projection in $\mathcal{C}(X, G)$ (and conversely). But on any compact set K , all but finitely many matrix coefficients of the $F_j|_K$ are zero. So by Lemma 5.11 the images of F_j and F_j^* are contained in a (standard) G -bundle. The theorem follows since $\mathcal{C}(X, G)$ is dense in $C^*(X, G)$. \square

6.8. THEOREM. *Let X be a G -compact proper G -space. The bijections in Lemma 6.5 determine a natural isomorphism between the Grothendieck group of G -bundles on X and the K_0 -group of the C^* -algebra $C^*(X, G)$.*

PROOF. If A is any C^* -algebra with an approximate unit consisting of projections, then the natural map from the Grothendieck group of projections in matrix algebras over A into $K_0(A)$ is an isomorphism. So the theorem follows from the previous result. \square

6.9. DEFINITION. If X is a G -compact proper G -space, and $j \in \mathbb{Z}/2\mathbb{Z}$, then denote by $K_G^j(X)$ the K_j -group of the C^* -algebra $C^*(X, G)$. If Y is a G -invariant closed subset of X , then denote by $K_G^j(X, Y)$ the K_0 -group of the ideal in $C^*(X, G)$ consisting of functions that vanish on Y .

By the above, $K_G^0(X)$ is the Grothendieck group of isomorphism classes of G -bundles on X .

The relative groups $K_G^j(X, Y)$ satisfy excision (of the strongest possible type, that $K_G^j(X, Y)$ depends only on $X \setminus Y$). Elementary K -theory provides functorial coboundary maps

$$\partial: K_G^j(Y) \longrightarrow K_G^{j+1}(X, Y),$$

and these give the groups $K_G^j(X, Y)$ the structure of a $\mathbb{Z}/2\mathbb{Z}$ -graded cohomology theory on G -compact proper G -spaces, in the sense that they fit into functorial long exact sequences

$$\cdots \longrightarrow K_G^j(X) \longrightarrow K_G^j(Y) \longrightarrow K_G^{j+1}(X, Y) \longrightarrow K_G^{j+1}(X) \longrightarrow \cdots$$

Although we have accessed this fact using C^* -algebra K -theory, this is also the main result of [LO01].

We conclude by reviewing the Gysin maps in equivariant K -theory that we shall need in the next section. Let E be a complex G -bundle over a G -compact proper G -manifold M . As we noted earlier, E carries a canonical Spin^c -structure. Form the sphere bundle \widehat{M} of the real bundle $\mathbb{R} \oplus E$, as in Section 3. The manifold M is equivariantly embedded as a retract in \widehat{M} using the section

$$M \ni m \mapsto (1, 0) \in \mathbb{R} \oplus E_m,$$

and associated to the embedding is a short exact sequence of K -theory groups

$$0 \longrightarrow K_G^0(\widehat{M}, M) \longrightarrow K_G^0(\widehat{M}) \longrightarrow K_G^0(M) \longrightarrow 0.$$

Let F be the complex G -bundle over \widehat{M} that we defined in Section 3, and denote by F_0 the complex G -bundle obtained by restricting F to M , then pulling back the restriction to \widehat{M} using the projection from \widehat{M} down to M . Because of the above exact sequence the difference $[F] - [F_0]$ defines an element of $K_G^0(\widehat{M}, M)$. The relative group is a module over $K_G^0(M)$ via pullback and tensor product, and multiplication against $[F] - [F_0]$ gives the *Thom homomorphism*

$$K_G^0(M) \longrightarrow K_G^0(\widehat{M}, M).$$

The complement $\widehat{M} \setminus M$ identifies with the total space of E via the map

$$E \ni e \mapsto \frac{1}{1+\|e\|^2} (1, e) \in \mathbb{R} \oplus E.$$

If $\iota: M \rightarrow N$ is an embedding of G -compact proper G -manifolds, and if the normal bundle to the embedding is identified with E , then the *Gysin map*

$$\iota_*: K_G^0(M) \rightarrow K_G^0(N)$$

is the composition

$$K_G^0(M) \xrightarrow{\text{Thom}} K_G^0(\widehat{M}, M) \xrightarrow{\cong} K_G^0(N, N') \longrightarrow K_G^0(N),$$

where $N' \subseteq N$ is the complement of a tubular neighborhood and the middle map is given by excision and the identification of the tubular neighborhood with E .

The Gysin map is functorial for compositions of embeddings. It is well-defined for embeddings of manifolds with boundary, as long as the embedding is transverse to the boundary of N , and carries the boundary of M into the boundary of N .

7. The Technical Theory

In this section we shall construct the homology groups $k_*^G(X)$ that were described in the introduction. They are obtained as direct limits of certain bordism groups.

7.1. DEFINITION. Let Z be a proper G -space and let E be a G -bundle on Z . A *stable (Z, E) -manifold* is a G -compact proper G -manifold M (possibly with boundary) together with an equivalence class of pairs (h, φ) , where:

- (a) $h: M \rightarrow Z$ is a continuous and G -equivariant map.
- (b) φ is an isomorphism of topological real G -bundles

$$\varphi: \mathbb{R}^r \oplus TM \longrightarrow \mathbb{R}^s \oplus h^*E,$$

for some $r, s \geq 0$. Here \mathbb{R}^r and \mathbb{R}^s denote the trivial bundles of ranks r and s (with trivial action of G on the fibers).

The equivalence relation is stable homotopy: (h_0, φ_0) and (h_1, φ_1) are equivalent if there is a homotopy $h: M \times [0, 1] \rightarrow Z$ between h_0 and h_1 and an isomorphism of real G -bundles over $M \times [0, 1]$

$$\varphi: \mathbb{R}^r \oplus TM \longrightarrow \mathbb{R}^s \oplus h^*E,$$

with $r \geq r_0, r_1$, that restricts to $I_{r-r_0} \oplus \varphi_0$ and $I_{r-r_1} \oplus \varphi_1$ at the two endpoints of $[0, 1]$.

If M is a stable (Z, E) -manifold with boundary, then its boundary may be equipped with a stable (Z, E) -structure by forming the composition

$$\mathbb{R}^r \oplus \mathbb{R} \oplus T\partial M \xrightarrow{\cong} \mathbb{R}^r \times TM|_{\partial M} \xrightarrow{\varphi} \mathbb{R}^s \oplus f^*E,$$

in which $\mathbb{R} \oplus T\partial M$ is identified with $TM|_{\partial M}$ by the ‘‘exterior normal first’’ convention.

7.2. DEFINITION. Let X be a G -finite proper G -CW-complex and let Y be a G -subcomplex of X . For $j = 0, 1, 2, \dots$ we define $\Omega_j^{(Z, E)}(X, Y)$ to be the group of equivalence classes of triples (M, α, f) , where

- (a) M is a smooth, proper, G -compact G -manifold of dimension j with a stable (Z, E) -structure.
- (b) α is a class in the group $K_G^0(M)$.
- (c) $f: M \rightarrow X$ is a continuous, G -equivariant map such that $f[\partial M] \subseteq Y$.

The equivalence relation is the obvious notion of bordism.

7.3. **REMARK.** Of course, the relation of bordism is arranged to incorporate the classes $\alpha \in K_G^0(M)$, so that if (M, α, f) is the boundary of (W, β, g) , then not only do we have that $M = \partial W$ and $f = g|_M$, but also the restriction map $K_G^0(W) \rightarrow K_G^0(M)$ takes β to α . One approach to the concept of bordism between manifolds with boundary is reviewed in [BHS07, Definition 5.5].

The sets $\Omega^{(Z,E)}(X, Y)$ are abelian groups. The group operation is given by disjoint union and the additive inverse of (M, α, f) is $(-M, \alpha, f)$. Here the *opposite* $-M$ is obtained by composing the bundle isomorphism φ with an orientation-reversing automorphism of \mathbb{R}^s . (See Lemma 8.3 for another description of the inverse.)

The groups $\Omega_j^{(Z,E)}(X, Y)$ constitute a homology theory on G -finite proper G -CW-complexes. Homotopy invariance follows from the bordism relation. The boundary maps

$$\partial: \Omega_j^{(Z,E)}(X, Y) \longrightarrow \Omega_{j-1}^{(Z,E)}(Y)$$

take (M, α, f) to $(\partial M, \alpha|_{\partial M}, f|_{\partial M})$. They fit into sequences

$$(7.1) \quad \cdots \longrightarrow \Omega_j^{(Z,E)}(X) \longrightarrow \Omega_j^{(Z,E)}(X, Y) \\ \longrightarrow \Omega_{j-1}^{(Z,E)}(Y) \longrightarrow \Omega_{j-1}^{(Z,E)}(X) \longrightarrow \cdots$$

whose exactness follows from direct manipulations with cycles, as follows. A cycle (M, α, f) for $\Omega_j^{(Z,E)}(Y)$, when mapped to $\Omega_j^{(Z,E)}(X, Y)$, represents the zero class since it is the boundary of $(M \times [0, 1], \alpha, f \circ \text{pr}_M)$. Conversely if a cycle for $\Omega_j^{(Z,E)}(X)$ is a boundary in $\Omega_f^{(Z,E)}(X, Y)$, then that boundary is a bordism from the given cycle to a cycle for $\Omega_j^{(Z,E)}(Y)$. So the sequence is exact at $\Omega_j^{(Z,E)}(X)$.

Exactness at $\Omega_{j-1}^{(Z,E)}(Y)$ is evident, as is the fact that the composition of the two maps through $\Omega_j^{(Z,E)}(X, Y)$ is zero. It remains to complete the proof of exactness at $\Omega_j^{(Z,E)}(X, Y)$. If a cycle (M, α, f) for $\Omega_j^{(Z,E)}(X, Y)$ maps to zero in $\Omega_{j-1}^{(Z,E)}(Y)$, then one can glue to (M, α, f) a null bordism for its image in $\Omega_{j-1}^{(Z,E)}(Y)$ to lift (M, α, f) to a cycle for $\Omega_j^{(Z,E)}(X)$. The glued manifold carries an evident stable (Z, E) -structure. As for the lifting of the K-theory class α , we use the fact that if α_1 and α_2 are equivariant K-theory classes on manifolds M_1 and M_2 that restrict to a common class on the a common boundary, then there is a K-theory class on $M_1 \cup M_2$ that restricts to α_1 and α_2 . This follows from the Mayer-Vietoris sequence

for equivariant K-theory, and therefore from the results of Lück and Oliver reviewed in Sections 5 and 6.

A map of pairs of G-CW-complexes $\phi: (X_1, Y_1) \rightarrow (X_2, Y_2)$ that is a homeomorphism from $X_1 \setminus Y_1$ to $X_2 \setminus Y_2$ induces an isomorphism on relative groups as follows. The open cells in $X_1 \setminus Y_1$ and $X_2 \setminus Y_2$ correspond to each other. By induction on the dimension of cells we can construct open G-invariant neighborhoods U_1 and U_2 of Y_1 and Y_2 , respectively, with G-equivariant deformation retractions $U_i \rightarrow Y_i$. Using the long exact sequence (7.1) we can replace Y_i with U_i . Given a cycle (M, α, f) for $\Omega_j^{(Z,E)}(X_2, U_2)$ we can, using a G-invariant collar of M , replace it by a cycle which omits Y_2 and therefore lifts to (X_1, U_1) . Similarly, we can achieve that a bordism for (X_2, U_2) of such normalized cycles avoids Y_2 and therefore lifts to (M_1, U_1) . Together, this implies that ϕ induces an isomorphism $\Omega_j^{(Z,E)}(X_1, Y_1) \rightarrow \Omega_j^{(Z,E)}(X_2, Y_2)$.

As they stand, the bordism groups $\Omega_j^{(Z,E)}(X, Y)$ are rather far from equivariant K-homology groups, most obviously because they are not 2-periodic. We shall obtain the technical groups $k_j^G(X, Y)$ by simultaneously forcing periodicity and removing dependence of the bordism groups on the pair (Z, E) .

Let M be a stable (Z, E) -manifold with structure maps h and φ , as in Definition 7.1 and let F be a complex hermitian G-bundle on Z of rank k . The pullback of F to M has a unique G-invariant smooth structure by Lemma 6.5, and so we may form the sphere bundle $S(\mathbb{R} \oplus h^*F)$, which is a G-compact proper G-manifold. It is also a stable $(Z, E \oplus F)$ -manifold. Indeed if $B(\mathbb{R} \oplus h^*F)$ is the unit ball bundle, and if p is the projection to M , then

$$TB(\mathbb{R} \oplus h^*F) \cong \mathbb{R} \oplus p^*T^*M \oplus h^*F$$

(once a complement to the vertical tangent bundle is chosen). So we obtain an isomorphism

$$\begin{aligned} \mathbb{R}^r \oplus TB(\mathbb{R} \oplus h^*F) &\cong \mathbb{R}^r \oplus \mathbb{R} \oplus p^*T^*M \oplus p^*h^*F \\ &\cong \mathbb{R}^s \oplus \mathbb{R} \oplus p^*h^*E \oplus p^*h^*F \end{aligned}$$

using the given stable (Z, E) -structure on M . We can then equip the sphere bundle with the stable $(Z, E \oplus F)$ -structure it inherits as the boundary of the ball bundle.

Suppose now that (M, α, f) is a cycle for the bordism group $\Omega_j^{(Z,E)}(X, Y)$. We can form from it the cycle $(\widehat{M}, \iota_!(\alpha), f \circ \pi)$ for the group $\Omega_{j+2k}^{(Z, E \oplus F)}(X, Y)$, where:

- (a) \widehat{M} is the sphere bundle for $\mathbb{R} \oplus h^*F$ with the stable $(Z, E \oplus F)$ -structure just described.

- (b) $\iota: M \rightarrow \widehat{M}$ is the inclusion of M into the sphere bundle given by the formula $m \mapsto (1, 0) \in \mathbb{R} \oplus F_{h(m)}$ and $\iota_*: K_G^0(M) \rightarrow K_G^0(\widehat{M})$ is the Gysin map, as described at the end of Section 6.
- (c) π is the projection from \widehat{M} to M .

Since this construction may also be carried out on bordisms between manifolds, we obtain a well-defined map on bordism classes.

7.4. DEFINITION. Let $k = \dim_{\mathbb{C}}(F)$. Denote by

$$\beta^F: \Omega_j^{(Z,E)}(X, Y) \longrightarrow \Omega_{j+2k}^{(Z,E \oplus F)}(X, Y)$$

the map determined by the above construction.

7.5. LEMMA. If F_1 and F_2 are G -bundles on Z of ranks k_1 and k_2 respectively, then

$$\beta^{F_2} \circ \beta^{F_1} = \beta^F: \Omega_j^{(Z,E)}(X, Y) \longrightarrow \Omega_{j+2k}^{(Z,E \oplus F)}(X, Y),$$

where $F = F_1 \oplus F_2$ and $k = k_1 + k_2$.

PROOF. Let $c = (M, \alpha, f)$ be a cycle for the bordism group $\Omega_j^{(Z,E)}(X, Y)$. The manifolds \widehat{M}_0 , \widehat{M}_1 and \widehat{M}_2 obtained from M by the modification processes underlying β^{F_1} , $\beta^{F_2} \circ \beta^{F_1}$ and β^F are fiber bundles over M whose fibers are the spheres in $\mathbb{R} \oplus F_1$ in the first case, the product of spheres in $\mathbb{R} \oplus F_1$ and $\mathbb{R} \oplus F_2$ in the second and the sphere in $\mathbb{R} \oplus F$ in the third. The product embeds as a hypersurface in the unit ball of $\mathbb{R} \oplus F$, for example via the map

$$((s_1, f_1), (s_2, f_2)) \mapsto \frac{1}{6}((2 + s_1)s_2, f_1, (2 + s_1)f_2),$$

and we obtain from this construction a bordism \widehat{W} between \widehat{M}_1 and \widehat{M}_2 . The map

$$j: t \mapsto (\frac{1}{2}(1 + t), 0, 0)$$

embeds $M \times [0, 1]$ into \widehat{W} , transversely to the boundary of \widehat{W} , and on the boundary components of $M \times [0, 1]$ the embedding restricts to the given embeddings of M into \widehat{M}_1 and \widehat{M}_2 . The embedding into \widehat{M}_1 is the composition of the embedding into \widehat{M}_0 with the embedding of \widehat{M}_0 into \widehat{M}_1 . The class $\alpha \in K_G^0(M)$ determines a class $\alpha \in K_G^0(M \times [0, 1])$ by homotopy invariance, and the triple $(\widehat{W}, j_!(\alpha), f \circ \pi)$ gives a bordism between the cycles representing $\beta^{F_2}(\beta^{F_1}(c))$ and $\beta^F(c)$, as required, using functoriality of the Gysin homomorphism to describe $\beta^{F_2}(\beta^{F_1}(c))$ in terms of the embedding $M \hookrightarrow \widehat{M}_1$. \square

Now fix a universal space $\underline{E}G$ as in Section 2. Let Z be a G -finite, G -subcomplex of $\underline{E}G$. In order to cope with the contingency that G might

be finite we shall modify the notion of G -standard bundle as advertised in Remark 5.7, so that standard G -bundles are now taken to be suitable subbundles of $Z \times \mathbb{C}[G_\infty]$.

Let E be a standard G -bundle over Z , as considered in Section 5. Form a partial order on the set of pairs (Z, E) by inclusion:

$$(Z_1, E_1) \leq (Z_2, E_2) \iff Z_1 \subseteq Z_2 \text{ and } E_1 \subseteq E_2|_{Z_1}.$$

According to the results of Section 5 this is a directed set.

7.6. DEFINITION. For $j \in \mathbb{Z}$ define groups $k_j^G(X, Y)$ to be the direct limits

$$k_j^G(X, Y) = \varinjlim_{(Z, E)} \Omega_{j+2\text{rank}(E)}^{(Z, E)}(X, Y)$$

over the directed set of all pairs (Z, E) , as above.

8. Proof of the Main Theorem

We aim to prove Theorem 4.6, that the geometric equivariant K -homology groups of Section 3 are isomorphic to the analytic groups of Section 4. We shall do so by comparing the technical groups of the previous section first to equivariant KK -theory and then to geometric K -homology.

The equivariant KK -groups determine a homology theory on G -finite proper G -CW pairs (or indeed on arbitrary second-countable G -compact proper G -CW pairs) if one defines the relative groups for a pair (X, Y) to be $KK_j^G(C_0(X \setminus Y), \mathbb{C})$. The boundary maps are provided by the boundary maps of the six-term exact sequence in KK -theory.

If (M, α, f) is a cycle for $\Omega^{(Z, E)}(X, Y)$, then an element of the Kasparov group $KK_j^G(C_0(X \setminus Y), \mathbb{C})$ may be defined as follows. Form the Dirac operator D on the interior of M using the Spin^c -structure associated to the given stable (Z, E) -structure on M . It determines a class

$$[D] \in KK_j^G(C_0(M \setminus \partial M), \mathbb{C}).$$

For example we may equip $M \setminus \partial M$ with a complete G -invariant Riemannian metric and then form a KK -class using $F = FD(I + D^2)^{-\frac{1}{2}}$ as in Section 4 (it does not depend on the choice of metric). Compare [HR00, Ch. 10], where the non-equivariant case is handled; the G -compact proper G -manifold situation is the same. We can then form the Kasparov product $\alpha \otimes [D] \in KK_j^G(C_0(M \setminus \partial M), \mathbb{C})$ and hence the class

$$f_*(\alpha \otimes [D]) \in KK_j^G(C_0(X \setminus Y), \mathbb{C})$$

more or less exactly as we did in Section 4.

8.1. THEOREM. *The correspondence that associates to a cycle (M, α, f) the element $f_*(\alpha \otimes [D])$ above is a natural transformation*

$$\mu: k_*^G(X, Y) \longrightarrow KK_*^G(C_0(X \setminus Y), \mathbb{C})$$

between homology theories. □

This is a mild elaboration of Theorem 4.5 and the equivariant counterpart of [BHS07, Theorem 6.1]. The equivariant case may be handled exactly as in [BHS07].

Our first goal is to show that this natural transformation is an isomorphism on G -finite proper G -CW-complexes. To do so it suffices to show that it is an isomorphism on the building blocks $X = G/H$ corresponding to finite subgroups H of G . The following observation clarifies what needs to be done in this case.

8.2. LEMMA. *Let H be a finite subgroup of G . There is a commutative diagram*

$$\begin{array}{ccc} k_*^G(G/H) & \xrightarrow{\mu} & KK_*^G(C_0(G/H), \mathbb{C}) \\ \text{Ind} \uparrow & & \uparrow \text{Ind} \\ k_*^H(\text{pt}) & \xrightarrow{\mu} & KK_*^H(\mathbb{C}, \mathbb{C}) \end{array}$$

in which the vertical maps are isomorphisms.

PROOF. The right-hand induction map is defined as follows. If \mathcal{H} is a Hilbert space, or Hilbert module, with unitary H -action, define $\text{Ind } \mathcal{H}$ to be the space of square-integrable sections of $G \times_H \mathcal{H}$. It carries a natural representation of $C_0(G/H)$, and if F is an H -equivariant Fredholm operator on \mathcal{H} , then operator $\text{Ind } F$ on $\text{Ind } \mathcal{H}$ given by the pointwise action of F determines a cycle for $KK_j^G(C_0(G/H), \mathbb{C})$.

The inverse to the induction map defined in this way is given by compression to the range of the projection operator determined by the indicator function of the identity coset in G/H (this function being viewed as an element of $C_0(G/H)$).

The left-hand induction map is defined in a similar fashion. We choose our model for $\underline{E}H$ to be a point, which we can include into $\underline{E}G$ as an H -fixed 0-cell, and we use the induced manifolds $\text{Ind } M = G \times_H M$, which map canonically to $G/H \subseteq \underline{E}G$. Note that any G -manifold that maps to G/H has this form. The construction of an inverse to induction and the proof that induction is an isomorphism are immediate upon noting that any G -map $f: \text{Ind } M \rightarrow \underline{E}G$ is G -equivariantly homotopic to one that factors through $G/H \subseteq \underline{E}G$. □

It suffices, therefore, to prove that the map

$$\mu: k_j^H(\text{pt}) \longrightarrow KK_j^H(\mathbb{C}, \mathbb{C})$$


is an isomorphism. The right hand group is isomorphic to the representation ring $R(H)$ when j is even and is zero when j is odd.

8.3. LEMMA. *Let (M, α, f) be a cycle for $k_j^G(X)$. If $\alpha = \alpha_1 + \alpha_2$ in $K_G^0(M)$, then*

$$[(M, \alpha, f)] = [(M, \alpha_1, f)] + [(M, \alpha_2, f)]$$

in $k_j^G(X)$.

PROOF. Suppose that $(M, \alpha, f) \in \Omega_{j+2k}^{(Z,E)}(X)$. Fix a bordism W between S^2 and $S^2 \sqcup S^2$ by situating two copies of the 2-sphere of radius $\frac{1}{4}$ inside the unit sphere. There are smooth paths I_1 and I_2 embedded into the bounding manifold W that connect the north and south poles of the large sphere to the south and north poles of the small spheres, that meet the spheres transversally, and that have trivial normal bundles in W .



The class $\alpha_1 \in K_G^0(M)$ pulls back to a class $\tilde{\alpha}_1 \in K_G^0(M \times I_1)$. Similarly, the class α_2 pulls back to a class $\tilde{\alpha}_2 \in K_G^0(M \times I_2)$. We obtain

$$\tilde{\alpha} := \tilde{\alpha}_1 \sqcup \tilde{\alpha}_2 \in K_G^0(M \times I) = K_G^0(M \times I_1) \oplus K_G^0(M \times I_2),$$

where $I = I_1 \sqcup I_2$. Now form the class

$$j_!(\tilde{\alpha}) \in K_G^0(M \times W),$$

where j is the inclusion of $M \times I$ into $M \times W$. If $\tilde{f}: M \times W \rightarrow X$ is the projection from $M \times W$ to M , followed by f , then $(M \times W, j_!(\tilde{\alpha}), \tilde{f})$ is a bordism between the images of (M, α, f) and $(M, \alpha_1, f) \sqcup (M, \alpha_2, f)$ under the map

$$\beta: \Omega_j(X)^{(Z,E)} \rightarrow \Omega_{j+2(k+1)}^{(Z,E \oplus \mathbb{C})}(X).$$

For this, observe that

$$\beta(M, \alpha, f) = (M \times S^2, i_!^N(\alpha_1 + \alpha_2), f) = (M \times S^2, i_!^N(\alpha_1) + i_!^S(\alpha_2), f)$$

where $i_N, i_S: M \rightarrow M \times S^2$ are the north pole inclusion and the south pole inclusion, which are G -homotopic, and the Gysin map is homotopy

invariant. Finally, $i_!^N(\alpha_1) = j_!(\tilde{\alpha}_1)|_{M \times S^2}$ by the compatibility of the Gysin map with restriction, so that $i_!^N(\alpha_1) + i_!^S(\alpha_2) = j_!(\tilde{\alpha})|_{M \times S^2}$.

In view of the definition of $k_j^G(X)$ the lemma is proved. \square

8.4. REMARK. The lemma shows that $-[(M, \alpha, f)] = [(M, -\alpha, f)]$ in $k_j^G(X)$.

8.5. PROPOSITION. *Let H be a finite group. The homomorphism*

$$\mu: k_*^H(\text{pt}) \longrightarrow \text{KK}_*^H(\mathbb{C}, \mathbb{C})$$

is an isomorphism.

PROOF. We shall prove that the homomorphism

$$\begin{aligned} R(H) &\longrightarrow k_0^H(\text{pt}) \\ \alpha &\mapsto (\text{pt}, \alpha, \text{Id}) \end{aligned}$$

is surjective, and that in addition, the group $k_1^H(\text{pt})$ is zero. This will suffice since $\text{KK}_0^H(\mathbb{C}, \mathbb{C}) \cong R(H)$, with the isomorphism being given by the above correspondence and the map μ , while $\text{KK}_1^H(\mathbb{C}, \mathbb{C}) = 0$.

Fix an element of $k_0^H(\text{pt})$ and represent it by a cycle (M, α) for $\Omega_{2n}^{(\text{pt}, E)}(\text{pt})$ (we drop the map $f: M \rightarrow \text{pt}$ from our notation here and below). Thus M is a $2n$ -dimensional Spin^c -manifold with a given stable isomorphism from its tangent bundle to a trivial bundle $E \times M$, where E is a complex representation of H :

$$\varphi: (\mathbb{R}^r \times M) \oplus TM \longrightarrow (\mathbb{R}^s \times M) \oplus (E \times M)$$

(for the purposes of the current argument it is convenient to vary our notation from that of the previous section and write trivial bundles as cartesian products). The manifold M may be equivariantly embedded in a finite-dimensional complex representation V of H . By composing with the vector space embedding of V into $\mathbb{R}^{r+s} \oplus E \oplus V$ as a summand we arrive at an H -equivariant embedding of the manifold M into $\mathbb{R}^{r+s} \oplus E \oplus V$. Write the trivial bundle $V \times M$ as the direct sum of TM and the normal bundle $N_V M$ of the embedding of M into V . The normal bundle of the embedding of M into $\mathbb{R}^{r+s} \oplus E \oplus V$ is then

$$N_{\mathbb{R}^{r+s} \oplus E \oplus V} M = (\mathbb{R}^{r+s} \times M) \oplus (E \times M) \oplus N_V M.$$

Using the vector bundle isomorphism φ we obtain isomorphisms

$$\begin{aligned} (\mathbb{R}^{r+s} \times M) \oplus (E \times M) \oplus N_V M & \\ &\cong (\mathbb{R}^r \times M) \oplus (\mathbb{R}^s \times M) \oplus (E \times M) \oplus N_V M \\ &\cong (\mathbb{R}^r \times M) \oplus (\mathbb{R}^r \times M) \oplus TM \oplus N_V M \\ &\cong (\mathbb{C}^r \times M) \oplus (V \times M). \end{aligned}$$

That is, the normal bundle $N_{\mathbb{R}^{r+s} \times E \times V}M$ is equipped with an isomorphism to the trivial bundle $W \times M$, where $W = \mathbb{C}^r \oplus V$.

Using the map

$$\beta^F: \Omega_{2n}^{(\text{pt}, E)}(\text{pt}) \longrightarrow \Omega_{2n+2k}^{(\text{pt}, E \oplus W)}(\text{pt})$$

we find that the given element of $k_0^H(\text{pt})$ is represented by the cycle $(\widehat{M}, \iota_*(\alpha))$. The manifold \widehat{M} is realizable as the boundary of a closed tubular neighborhood X of M in $E \oplus W \oplus \mathbb{R}$. By enclosing X in a large ball we construct a bordism Y between \widehat{M} and a sphere $S \subseteq V \oplus W \oplus \mathbb{R}$. The union of X and Y is the ball bounded by the sphere S , and by applying the Mayer-Vietoris sequence in H -equivariant K -theory to the decomposition $X \cup Y$ of the ball, we find that the class $\iota_*(\alpha) \in K_H^0(\widehat{M})$ may be written as a sum $b_X + b_Y$, where b_X is the restriction of a class in $K_H^0(X)$ and b_Y is the restriction of a class in $K_H^0(Y)$. By the previous lemma,

$$[(\widehat{M}, b)] = [(\widehat{M}, b_X)] + [(\widehat{M}, b_Y)].$$

The first class is zero in $\Omega_{2n+2k}^{(\text{pt}, E \oplus W)}(\text{pt})$, while the second is equal to some class $[(S, c)]$ thanks to the bordism Y . We have therefore shown that every class in $k_0^H(\text{pt})$ represented by a cycle (S, c) , where S is the sphere in some $V \oplus W \oplus \mathbb{R}$.

To complete the proof we invoke Bott periodicity. The class c is a sum $c_0 + c_1$ where c_0 is represented by a trivial bundle (one that extends over the ball) and c_1 is in the image of the Gysin map associated to the inclusion

$$\text{pt} \mapsto (0, 0, 1) \in S \subseteq V \oplus W \oplus \mathbb{R}.$$

This completes the computation of $k_0^H(\text{pt})$.

The proof that $k_1^H(\text{pt}) = 0$ is essentially the same. Every cycle is equivalent to one of the form $[(S, c)]$, where S is the sphere in a complex representation of H . But the sphere is odd-dimensional and by Bott periodicity every class in $K_H^0(S)$ extends over the ball. So (S, c) is a boundary. \square

8.6. COROLLARY. *If X is a G -finite proper G -CW-complex, then the map*

$$\mu: k_*^G(X) \longrightarrow \text{KK}_*^G(C_0(X), \mathbb{C})$$

is an isomorphism. \square

Next we need to relate the technical groups $k_j^G(X)$ to the geometric groups $K_j^G(X)$.

8.7. LEMMA. *The class in $K_j^G(X)$ of an equivariant K -cycle (M, E, f) depends only on M , f and the class of the G -bundle E in the Grothendieck group $K_G^0(X)$.*

PROOF. Fixing M and f , the map that associates to a G -bundle E on M the class of (M, E, f) in $K_j^G(X)$ is additive, and so extends to a map from the Grothendieck group $K_G^0(X)$ into $K_j^G(X)$. \square

Thanks to the lemma, we can attach a meaning to the class in the geometric group $K_j^G(X)$ of any triple (M, α, f) , whenever M is a G -compact proper G -Spin^c-manifold, f is an equivariant map from M to X , and $\alpha \in K_G(M)$. In particular, we can do so when M is a stable (Z, E) -manifold. We obtain in this way a natural transformation

$$\Omega_j^{(Z, E)}(X) \longrightarrow K_j^G(X)$$

8.8. LEMMA. *If F is any G -bundle over Z of rank k , then the diagram*

$$\begin{array}{ccc} \Omega_j^{(Z, E)}(X) & \longrightarrow & K_j^G(X) \\ \beta^F \downarrow & & \parallel \\ \Omega_{j+2k}^{(Z, E \oplus F)}(X) & \longrightarrow & K_{j+2k}^G(X) \end{array}$$

is commutative.

PROOF. This follows from the definitions of the Gysin homomorphism and vector bundle modification. \square

We obtain therefore a natural transformation

$$k_j^G(X) \longrightarrow K_j^G(X)$$

that fits into a commuting diagram

$$\begin{array}{ccc} k_j^G(X) & \longrightarrow & K_j^G(X) \\ \parallel & & \downarrow \mu \\ k_j^G(X) & \xrightarrow{\mu} & KK_j^G(C_0(X), \mathbb{C}). \end{array}$$

8.9. LEMMA. *For every j the map from $k_j^G(X)$ into the equivariant geometric K-homology group $K_j^G(X)$ is surjective.*

PROOF. Let (M, α, f) be an equivariant K-cycle for X . The manifold M maps equivariantly to a G -finite subcomplex Z of the universal space $\underline{E}G$ via h .

The tangent bundle for M (indeed its complexification) embeds as a summand of a G -bundle that is pulled back from a standard G -bundle on Z via h : $M \rightarrow Z$ (see Corollary 5.2 and the comment following it). Thus there is an isomorphism of real bundles

$$TM \oplus F_0 \cong h^*E$$

where F_0 is a real G -bundle on M and E is a complex G -bundle on Z . By adding a trivial bundle if necessary, we obtain an isomorphism

$$TM \oplus F_1 \cong h^*E \oplus \mathbb{R}^s,$$

where $F_1 = F_0 \oplus \mathbb{R}^s$ has *even* fiber dimension. By the two out of three principle for Spin^c -structures from Section 3, the bundle F_1 carries a Spin^c -structure whose direct sum with the given Spin^c -structure on TM is the direct sum of the Spin^c -structure on h^*E associated with its complex structure and the trivial Spin^c -structure on \mathbb{R} . If we carry out a vector bundle modification using F_1 , then we obtain an equivariant K -cycle $(M, E, f)^\wedge$ that is equivalent to (M, E, f) and for which \widehat{M} carries a stable (Z, E) -structure compatible with its Spin^c -structure. \square

With this, as we pointed out in the introduction, the proof of Theorem 4.6 is complete.

9. The Baum-Connes Assembly Map

Let G be a countable discrete group. The essence of the Baum-Connes conjecture for G is the assertion that every class in the K -theory of the reduced group C^* -algebra $C_r^*(G)$ arises as the index of an elliptic operator on a G -compact proper G -manifold, and that in addition the only relations among these indexes arise from geometric relations (such as for example bordism) between the operators. The conjecture arose from a K -theoretic analysis of Lie groups and of crossed product algebras related to foliations. However here we shall discuss only the C^* -algebras of discrete groups.

To make their conjecture precise, Baum and Connes constructed in [BC00] geometric groups $K^j(G)$ from cycles related to the symbols of equivariant elliptic pseudodifferential operators, and an equivalence relation related to the Gysin map in K -theory. They then defined an index map

$$\mu: K^j(G) \longrightarrow K_j(C_r^*(G))$$

that they conjectured to be an isomorphism.

Although the Dirac operator on a Spin^c -manifold did not play a central role in [BC00], it is nonetheless a fairly routine matter to identify the geometric group defined there with the group generated from cycles (M, E) , where:

- (a) M is a G -compact proper G - Spin^c -manifold, all of whose components have either even or odd dimension, according as j is 0 or 1, and
- (b) E is a complex G -bundle on M .

The equivalence relation between cycles is generated by bordism, direct sum/disjoint union and vector bundle modification, exactly as in Section 3, except that here there is no reference space X , nor any map from M to X .

Compare [Bau04], where the conjecture for countable discrete groups is formulated in precisely this way.

It follows from the universal property of the space $\underline{E}G$ that there is an isomorphism

$$K^j(G) \cong \varinjlim_{X \subseteq \underline{E}G} K_j^G(X),$$

where on the right is the direct limit of the geometric K-homology groups of the G-finite subcomplexes of the G-CW-complex $\underline{E}G$.

In a later paper [BCH94], Baum, Connes and Higson defined an assembly map

$$\mu: \varinjlim_{X \subseteq \underline{E}G} KK_j^G(C(X), \mathbb{C}) \longrightarrow K_j(C_r^*(G))$$

Its relation to the original Baum-Connes map is summarized by the commutative diagram

$$\begin{array}{ccc} K^j(G) & \xrightarrow{\cong} & c\varinjlim_{X \subseteq \underline{E}G} K_j^G(X) & \xrightarrow{\mu} & \varinjlim_{X \subseteq \underline{E}G} KK_j^G(C(X), \mathbb{C}) \\ \mu \downarrow & & & & \downarrow \mu \\ K_j(C_r^*(G)) & \xrightarrow{=} & & & K_j(C_r^*(G)), \end{array}$$

where the horizontal map labelled (yet again) μ is the one analyzed in this paper, and shown to be an isomorphism. Because it is an isomorphism, the reformulation of the Baum-Connes conjecture in [BCH94] is equivalent to the original in [BC00] for discrete groups.

Despite the discovery some years ago of counterexamples to various extensions of the Baum-Connes conjecture (see [HLS02]), there is, as of today, no known counterexample to the conjecture as reviewed here.

References

- [Bau04] P. Baum. On the index of equivariant elliptic operators. In *Operator algebras, quantization, and noncommutative geometry*, volume 365 of *Contemp. Math.*, pages 41–49. Amer. Math. Soc., Providence, RI, 2004.
- [BC00] P. Baum and A. Connes. Geometric K-theory for Lie groups and foliations. *Enseign. Math. (2)*, 46(1-2):3–42, 2000.
- [BCH94] P. Baum, A. Connes, and N. Higson. Classifying space for proper actions and K-theory of group C^* -algebras. In *C^* -algebras: 1943–1993 (San Antonio, TX, 1993)*, volume 167 of *Contemp. Math.*, pages 240–291. Amer. Math. Soc., Providence, RI, 1994.
- [BD82] P. Baum and R. G. Douglas. K homology and index theory. In *Operator algebras and applications, Part I (Kingston, Ont., 1980)*, volume 38 of *Proc. Sympos. Pure Math.*, pages 117–173. Amer. Math. Soc., Providence, R.I., 1982.
- [BHS07] P. Baum, N. Higson, and T. Schick. On the equivalence of geometric and analytic K-homology. *Pure Appl. Math. Q.*, 3(1, part 3):1–24, 2007.

- [Bla98] B. Blackadar. *K-theory for operator algebras*, volume 5 of *Mathematical Sciences Research Institute Publications*. Cambridge University Press, Cambridge, second edition, 1998.
- [BOOS09] P. Baum, H. Oyono-Oyono, and T. Schick. Equivariant geometric K-homology for compact Lie group actions, 2009. arXiv.org:0902.0641.
- [EM08] H. Emerson and R. Meyer. Bivariant K-theory via correspondences, 2008. arXiv.org:0812.4949.
- [HLS02] N. Higson, V. Lafforgue, and G. Skandalis. Counterexamples to the Baum-Connes conjecture. *Geom. Funct. Anal.*, 12(2):330–354, 2002.
- [HR00] N. Higson and J. Roe. *Analytic K-homology*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000. Oxford Science Publications.
- [Kas88] G. G. Kasparov. Equivariant KK-theory and the Novikov conjecture. *Invent. Math.*, 91(1):147–201, 1988.
- [LO01] W. Lück and R. Oliver. The completion theorem in K-theory for proper actions of a discrete group. *Topology*, 40(3):585–616, 2001.
- [Lüc05] W. Lück. Survey on classifying spaces for families of subgroups. In *Infinite groups: geometric, combinatorial and dynamical aspects*, volume 248 of *Progr. Math.*, pages 269–322. Birkhäuser, Basel, 2005.

P.B.: Department of Mathematics, Penn State University, University Park, PA 16802. Email: baum@math.psu.edu.

N.H.: Department of Mathematics, Penn State University, University Park, PA 16802. Email: higson@math.psu.edu.

T.S.: Mathematisches Institut, Universität Göttingen, Bunsenstr. 3, D-37073 Göttingen, Germany. Email: thomas.schick@uni-math.gwdg.de.